




10/540160
PCT/GB 2003 / 0 D 5 5 4 4

18 DECEMBER 2003

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




P. Mahoney

Signed

Dated 3 December 2003

BEST AVAILABLE COPY

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



The
Patent
Office

28DEC02 773531-6 06:06
P01/7700 0.00-0230095.2

1/77

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet, from the Patent Office to help you fill in this form)

1.	Your reference	CDK2066		
2.	Patent application number (The Patent Office will fill in this part)	0230095.2		
3.	Full name, address and postcode of the or of each applicant (underline all surnames)	RHODIA CONSUMER SPECIALTIES LIMITED Oak House Reeds Crescent Watford Hertfordshire WD24 4QP. 8486797001		
	Patents ADP number (if you know it)			
	If the applicant is a corporate body, give the country/state of its incorporation			
4.	Title of the invention	PHOSPHORODIAMIDITE		
5.	Name of your agent (if you have one)	Barker Brettell		
	"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	138 Hagley Road Edgbaston Birmingham B16 9PW		
	Patents ADP number (if you know it)	7442494002		
6.	If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of Filing (day/month/year)
7.	If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day/month/year)	
8.	Is a statement of inventorship and of right to grant of a patent required in support of this request (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))	YES		

Patents Form 1/77

Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form -

Description 5 + 5

Claim(s) 2 + 2

Abstract -

Drawing(s) -

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*) -

Request for preliminary examination 1 /
(*Patents Form 9/77*)

Request for substantive examination -
(*Patents Form 10/77*)

Any other documents -
(*please specify*)

11. I/We request the grant of a patent on the basis of this application.

Signature
Barker Brettell
Barker Brettell

Date
23 December 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Colin D. Kinton

Tel: 0121 456 1364

Warning
After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes
a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 01645 500505
b) Write your answers in capital letters using black ink or you may type them.

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.

e) Once you have filled in the form you must remember to sign and date it.

f) For details of the fee and ways to pay please contact the Patent Office.

PHOSPHORODIAMIDITE

The present invention relates to an improved method for the production of phosphorodiamidites, phosphorodiamidites produced by way of such a method and the use of such phosphorodiamidites.

Production of phosphorodiamidites has become increasingly important in the biotechnology industry. Phosphorodiamidites are used as intermediates in the manufacture of novel anti-neoplastic agents.

To be suitable for use in such industries phosphorodiamidites must be of high purity. Such phosphorodiamidites must also contain low levels of bis-(2-cyanoethyl) phosphorodiamidite (the 'diester').

This impurity is known to be a significant by product in the synthesis of 2-cyanoethyl tetraisopropylphosphorodiamidite, a commercially important intermediate in the synthesis of oligonucleotides.

As phosphorodiamidites are very air sensitive and thermally unstable, their purification is, at present, complex and expensive. Hitherto, known processes of extraction and purification of phosphorodiamidites often involve multi-stage synthetic procedures which demand the chemical isolation of intermediate materials and require extensive purification procedures prior to the isolation of high purity phosphorodiamidite products.

The present invention aims to ameliorate the aforementioned disadvantages of phosphorodiamidite production.

Accordingly, the present invention provides a method of phosphorodiamidite production which method comprises the steps of

reacting a phosphorus trihalide with a dialkyl amine in a polar solvent to form an intermediate compound and subsequently reacting the intermediate compound with a hydroxyalkyl compound and a dialkyl amine, in the presence of a non-polar co-solvent.

5

Following filtration to remove the solid by-product, the two solvents form separate layers. This is advantageous as the upper, non-polar solvent, layer contains the high-purity phosphorodiamidite product. The lower, polar solvent, layer contains impure product contaminated with diester and other unwanted by-products. The upper layer is then subjected to vacuum-stripping to remove the solvent, leaving the desired product with greater than 96% purity and containing less than 1% of the diester impurity. The yield of the product can further be increased by optionally rewashing the polar solvent layer with a further quantity of non-polar solvent, to give non-polar solvent solution containing pure product, from which can then be isolated high-purity phosphorodiamidite.

15

Advantageously, impure product contaminated with diester and other impurities which would otherwise be unsuitable for commercial use can be extracted and purified by use of the solvent purification procedure. Phosphorodiamidite products are preferentially soluble in the non-polar co-solvent whereas the diester and other unwanted polar by-products are insoluble and remain in the polar solvent layer.

20

Preferably, the phosphorus trihalide is phosphorus trichloride. Alternatively, the phosphorus trihalide is phosphorus tribromide.

25

The dialkyl amine is preferably diisopropylamine. Alternatively the dialkyl amine may be dimethylamine, diethylamine, di-n-propylamine, di-n-butylamine, di-isobutylamine or di-tert-butylamine.

30

The polar solvent is preferably a nitrile compound, in particular, acetonitrile. Alternatively the polar solvent may be propionitrile or benzonitrile.

- 5 The hydroxyalkyl compound is preferably hydroxypropionitrile. Alternatively the hydroxyalkyl compound may be methanol, tert-butyl alcohol or other suitable hydroxyalkyl compounds which are known to be suitable for the manufacture of phosphorodiamidites.
- 10 The alkane co-solvent is preferably heptane or hexane. Other suitable C₅ to C₉ aliphatic hydrocarbons include pentane. Suitable alicyclic hydrocarbons include, for example, cyclohexane.

The ratio of polar solvent to non-polar solvent is suitably around 1:1.

- 15 The method according to the invention provides a phosphorodiamidite compound according to Formula I:



- 20 wherein R is a C₁ to C₄ alkyl, hydroxyalkyl or oxyalkyl group; and n is a whole number of from 1 to 4.

The compound according to formula I is preferably 2-cyanoethyl tetraisopropyl phosphorodiamidite wherein R is isopropyl, and n = 2.

25

The present invention also provides the use of a compound of formula I in the synthesis of oligonucleotides.

- 30 The present invention will now be illustrated, merely by way of example, as follows:

Example 1**Manufacture of 2-cyanoethyl tetraisopropyl phosphorodiamidite using hexane co-solvent**

5

27.5g of phosphorus trichloride at ambient temperature was added to a stirred mixture of acetonitrile (200g) and diisopropylamine (121g) over 1 hour. 200g of hexane is then added followed by 14g of hydroxypropionitrile at ambient temperature over 30 minutes. The reaction mixture is then stirred for 1 hour and is then filtered to remove the solid by-product. The upper hexane layer of the filtered reaction mixture is separated and subjected to vacuum stripping to remove the hexane solvent. This leaves 20g of 2-cyanoethyl tetraisopropylphosphorodiamidite which has a purity of 96.9% when analysed by ^{31}P -NMR. The lower acetonitrile layer is stirred with a further 200g of hexane for 2 hours. The upper hexane layer from this re-extraction contains product of 98% purity when assayed by ^{31}P -NMR. Following vacuum stripping a further 11g of high purity 2-cyanoethyl tetraisopropylphosphorodiamidite is isolated.

20

Example 2**Manufacture of 2-cyanoethyl tetraisopropylphosphorodiamidite using heptane co-solvent**

25

27.5g of phosphorus trichloride was added to a stirred mixture of 200g of acetonitrile and 121g of diisopropylamine at ambient temperature. 200g of heptane was then added to this mixture followed by 14.3g of hydroxypropionitrile at ambient temperature over 30 minutes. The reaction mixture was then stirred for an hour and was then filtered to remove the solid by-product. The upper heptane layer was then separated

30

and subjected to vacuum stripping to remove the heptane solvent leaving 22g of 2-cyanoethyl tetraisopropylphosphorodiamidite which had a purity of 96.7% when assayed by ^{31}P -NMR.

5 Example 3

Purification of low purity 2-cyanoethyl tetraisopropylphosphorodiamidite

- 10 60g of low purity 2-cyanoethyl tetraisopropylphosphorodiamidite (92%
purity when assayed by ^{31}P -NMR containing 1.3% diester) was added to a
mixture of 200g acetonitrile and 200g of heptane after stirring for ten
minutes the upper heptane layer was separated and the lower acetonitrile
layer stirred with a further 200g of heptane for a further 10 minutes. The
15 second heptane fraction was then separated and the two heptane fraction
subsequently combined and subjected to vacuum stripping to remove
heptane solvent. 30g of 2-cyanoethyl tetraisopropylphosphorodiamidite
was obtained at a purity of 98.3% when assayed by ^{31}P -NMR. This
extracted phosphorodiamidite compound contained less than 0.1% of the
20 diester impurity.

CLAIMS

1. A method of phosphorodiamidite production which method comprises the steps of reacting a phosphorus trihalide with a dialkyl
5 amine in a polar solvent to form an intermediate compound and subsequently reacting the intermediate compound with a hydroxyalkyl compound and a dialkyl amine, in the presence of a non-polar co-solvent.
2. A method as claimed in Claim 1 in which the phosphorus trihalide
10 is phosphorus trichloride.
3. A method as claimed in Claim 1 in which the phosphorus trihalide is phosphorus tribromide.
- 15 4. A method according to any one of Claims 1 to 3 in which the dialkyl amine is diisopropylamine.
5. A method as claimed in any one of Claims 1 to 3 in which the
20 dialkyl amine is selected from the group consisting of dimethylamine, diethylamine, di-n-propylamine, di-n-butylamine, di-isobutylamine or di-tert-butylamine.
6. A method as claimed in any one of the preceding claims in which
25 the polar solvent is a nitrile compound.
7. A method as claimed in Claim 6 in which the nitrile compound is acetonitrile.
8. A method as claimed in Claim 6 in which the polar solvent is
30 propionitrile or benzonitrile.

9. A method as claimed in any one of the preceding claims in which the hydroxyalkyl compound is hydroxypropionitrile.
10. A method as claimed in any one of Claims 1 to 8 in which the hydroxyalkyl compound is methanol or tert-butyl alcohol.
11. A method as claimed in any one of Claims 1 to 10 in which the alkane co-solvent is a C₅ to C₉ aliphatic hydrocarbon.
12. A method as claimed in any one of Claims 1 to 10 in which the alkane co-solvent is an alicyclic hydrocarbon.
13. A method according to any one of the preceding claims in which the ratio of polar solvent to non-polar solvent is 1:1.
14. A phosphorodiamidite compound produced by the method of any one of Claims 1 to 13 and having the General Formula (I):
- $$(R_2 N)_2-P-O(CH_2)_n-CN \quad (I)$$
- wherein R is a C₁ to C₄ alkyl, hydroxyalkyl or oxyalkyl group; and n is a whole number of from 1 to 4.
15. A compound according to Claim 14 which is 2-cyanoethyl tetraisopropyl phosphorodiamidite.
16. The use of a compound as claimed in Claim 14 or Claim 15 in the synthesis of oligonucleotides.
17. A phosphorodiamidite compound, substantially as hereinbefore described with reference to the Examples.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.